



OPEN

Ecological and historical factors behind the spatial structure of the historical field patterns in the Czech Republic

Václav Fanta^{1✉}, Jaromír Beneš^{2,3}, Jan Zouhar^{1,4}, Volha Rakava¹, Ivana Šitnerová^{2,3}, Kristina Janečková Molnárová¹, Ladislav Šmejda^{1,5} & Petr Sklenicka¹

Historical field systems are an essential part of the traditional cultural landscape of societies with primarily agricultural subsistence. They embody many functions and values, as they affect the productional, ecological and hydrological functioning of the landscape, its cultural values, the way people perceive the landscape, and their impact on present-day farming. As an aspect of the historical landscape, field systems are a topic investigated in landscape archaeology, environmental studies, historical geography, landscape ecology, and related disciplines. Historical field systems can form many complex spatial structures, shapes and patterns. This paper focuses on identifying environmental and historical/cultural driving forces during the formation and the historical development of various field pattern types. We worked with 523 settlements established in the medieval to the early modern period (approx. 900–1600 AD) in the present-day Czech Republic. We have determined the proportions of different field pattern types in the examined cadastres and have statistically compared them with a variety of environmental and geographical predictors. Our results indicate a strong influence of environmental predictors (terrain undulation, cadastre size), the impact of specific historical events and associated social changes (e.g. land confiscations by the state in the seventeenth century), and a significant relationship between field pattern types and settlement layout types. Furthermore, we have observed the different adaptations of field pattern types to similar environmental conditions, as well as the impact of social and political factors on the processes of landscape formation. Our paper provides the first detailed analysis of the geographical distribution of traditional field systems on the scale of an entire modern state, and emphasizes the importance of transdisciplinary research on cultural landscapes.

General introduction. Historical field systems and their remnants are an essential part of the historical cultural landscape. As a typical representation of the “combined work of nature and of man”¹, their pattern constitutes the image of the organically evolved cultural landscape and forms an integral part of a country’s cultural heritage^{2–5}. The history of field systems also significantly impacts the ecological function of present-day farming landscapes. Elements of these systems, especially historical field margins, control the physical, chemical and biological fluxes in the landscape⁶, mitigate soil erosion⁷, stabilize the landscape’s hydrological regime^{8,9}, reduce pesticide drift and fertilizer misplacement^{10,11}, and serve as a buffer against nitrates and for water protection¹². Furthermore, the remnants of historical field systems contribute to biodiversity conservation, serving as a habitat and a conduit for wildlife^{2,7,13,14}. From the standpoint of landscape ecology, the remnants of traditional field systems form specific landscape features¹⁵, which incorporate the spatial structure of the landscape, historical works of man, archaeological complexity and various ecological functions^{6,16,17}. We also want to stress that the

¹Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, 165 00 Prague – Suchbát, Czech Republic. ²Institute of Archaeology, Faculty of Philosophy, University of South Bohemia, Branišovská 31, 370 05 České Budějovice, Czech Republic. ³Laboratory of Archaeobotany and Palaeoecology, Faculty of Science, University of South Bohemia, Na Zlaté stoce 3, 370 05 České Budějovice, Czech Republic. ⁴Prague University of Economics and Business, W. Churchill Sq. 1938/4, 130 67 Prague 3 – Žižkov, Czech Republic. ⁵Department of Anthropology, Faculty of Arts, University of West Bohemia, Univerzitní 8, 306 14 Plzeň, Czech Republic. ✉email: fanta.vaclav@gmail.com

ecological functions of historical field systems and hedgerows have dramatically changed since the end of traditional agriculture¹⁸. Last but not least, the historical field systems as key macrostructures also form the image of the landscape and its aesthetic values^{7,19}.

Historical field systems, and knowledge gained from studying them, also play a significant role in modern sustainable agriculture and landscape planning^{20,21}. For example, with the ever-increasing impact of climate change, the hydrological function of remnants of historical field systems and their ability to mitigate erosion and retain water have become critical (cf.^{22,23}). In addition, the historical field pattern reflects the land ownership structure¹⁵ and social relationships between members of historical society²⁴, which makes it an essential topic for historians, archaeologists and historical geographers^{25–30}. In terms of spatial archaeology, historical field systems form one of the largest landscape macrostructures (or landscape “superartifacts” sensu Chang³¹), retaining information about their chronology, function, and environmental characteristics^{32–34}. The historical field pattern also co-creates a country’s cultural identity^{35,36}. Unfortunately, these landscape structures have tended to disappear or to have been abandoned from the middle of the twentieth century onwards^{37,38}.

To sum up, historical field systems and their remnants contribute with many values to the present-day cultural landscape: (a) they take their part in forming the “image of the landscape”, (b) they affect its ecological and hydrological functions, (c) they are historical witnesses of ancient times, and they reflect agriculture in the past and land-ownership traditions, and (d) they have a strong influence on present-day farming. In brief, the remnants of historical field systems can be remarkable from many points of view. This paper will focus on various types of spatial arrangement of historical field patterns in the Czech Republic, their driving forces and their spatial distribution.

Primary forms of historical field systems. Transformation of the landscape into organized field systems has taken place in many parts of the world^{39,40}. The most significant form of historical field systems are well visible *terraced fields*. This phenomenon is probably related to the origin of historical state organization, and it prevailed in agrarian usage of foothills and mountainous areas with more prominent dynamics of relief⁴¹. Terracing is an ancient technology; the oldest known (proto)terraces (*paddy fields*) were already constructed in the Neolithic period in China⁴². The oldest evidence of terraces is recorded from the Levant around 6000 BC^{43,44}. At a Neolithic site in Croatia, excavation of terraces associated with the site has been reported⁴⁵. The non-agricultural terraces were also used in the famous Neolithic site of Lepenski Vir in Serbia⁴⁶. Some studies admit the existence of terraces in the Aegean area in the Bronze Age⁴⁷. Further evidence of terraces is known from South-East Asia around 3000 BC⁴⁶. The oldest examples outside the Mediterranean area and South-East Asia have been reported from the Andes around 2400 BC⁴⁸. In the Alpine Region and the Maya Lowlands, the oldest terracing practices date back to the Iron Age⁴⁹. All these examples correlate well with the introduction of agriculture.

In addition to their agricultural function, terraced fields help retain water and prevent soil erosion^{15,49,50}. Terraced fields are found on all continents. They are present in Asia⁵¹, South America⁵², Central and North America⁵³, and the mountainous regions of Africa⁵⁴ and the Near East³². In Europe, terraced fields are located mainly in the Mediterranean area^{33,55}, in the Alpine region⁵⁶, and in Central Europe^{15,53}. There are several types of terraces, but all are built with two elements—the terraced platform and the terraced slope. A traditional terraced slope is a dry stone wall structure, or a slope reinforced with stones grubbed out during farmland cultivation and later covered with earth and grassed over⁵⁷. Modern terraces are usually grassed slopes. Terraces are typically used as an agricultural space, frequently for growing grapes and other fruit^{57–59}, olives⁶⁰, vineyards⁵⁸ and for grazing⁵⁸. These terraces are typical, for example, in France⁶¹, Slovenia⁵⁷, Portugal⁶², Spain⁶³ and Greece⁶⁴.

The second significant form of the historical field system is characterized by visible boundaries of the parcels³⁹. These field systems are called *bocage* or *hedgerow patterns* in western Europe⁶, *Flur* in Germany^{65–68}, and *plužina* in the Czech Republic². The boundaries between parcels were mostly marked by small walls, banks, steep slopes, pathways, or by lines of trees or bushes. These landscape elements also provided important secondary functions, e.g. they affected the surface water flow or served as a source of firewood¹⁵. In the present-day landscape, these traditional agricultural field patterns mostly appear as mosaics of small-scale arable fields, pastures, orchards or vineyards. Typically, the fields in these landscapes are defined by hedgerows composed of trees and shrubs, often unmanaged or with a low level of management (cf.^{13,69}). In some parts, there can be small woodland patches for obtaining timber. This landscape structure results from the historical development of agriculture, and it constitutes a traditional agroforestry system, which is typical for Central Europe^{15,70,71}.

The third significant form of the historical field system is so-called *open fields*, well-known from Great Britain^{17,30,72}. In these extensive systems, individual parcels are not separated by walls or other boundaries¹⁵. Renes⁷³ explains that the long and narrow shapes of these fields were generated using a heavy plough, which allowed only limited manipulation. This agrarian landscape setting following older prehistoric field systems⁷⁴ probably appeared in the tenth century and lasted until the sixteenth century^{15,73}.

Historical field pattern typologies. Several typological schemes have differentiated these field systems according to their spatial division, shape, size, terrain, etc. The individual national/regional typologies reflect the types present in the respective country. Authors from the Mediterranean region usually describe types of terraced fields. In contrast, Central European authors focus on field systems with visible boundaries (e.g. *Flur* in Germany, *plužina* in the Czech Republic). In some cases, these typologies may overlap.

There are two ways to differentiate agricultural terraces: horizontal layout (planform) and vertical profile. The horizontal layout can strictly follow the contour lines, can form braided structures or lynchets⁷⁵, or, alternatively, they can compose wave-like terraces⁴⁹ or off-contour terraces⁷⁶. Moreover, Brown et al.⁷⁵ identified regular box-like terrace structures and, together with Wei et al.⁴⁹, referenced pocket/half-moon shaped terraces. Pocket terraces also have a wall-free “spontaneous” version⁷⁶. Based on the vertical profile, authors have identified

four⁷⁷, six⁷⁵ or seven types⁴⁹ of terraces. The divisions are primarily based on the construction material⁷⁷ or on the terrain gradient of the levelled part of the terraces^{49,75,77}. The levelled part can also have a reversed slope or can include a ditch to increase the water retention capacity⁷⁸.

Unlike terraced fields, field systems with visible boundaries exhibit great diversity in their shapes, as they are constructed in less steep terrain. These field systems can be divided into many types, according to the spatial division of land ownership (individual homesteads or noble estates) and the terrain where they were built⁷⁹. There are several typologies of these plužina field systems: Löw and Michal⁷⁹ identified six types of plužina field pattern, Láznicka⁸⁰ identified seven types, while the broadest classification was introduced by Černý⁸¹, who divided the plužina field system into nine types. These typologies differ in details and in terminology. However, all of them recognize the following basic types: (a) an irregular, blocky structure, often called segmental or blocky plužina (Fig. 1A), (b) narrow and medium-length strips parallel to each other, arranged into large irregular blocks, called sectional plužina (Fig. 1C), and (c) very long parallel strips adjacent to each homestead, called croft or longitudinal plužina (Fig. 1D). Other types of field patterns with visible boundaries have also been reported from the Mediterranean landscape, e.g. more or less regular flat fields and sloping fields, which occur mostly in plain terrain with fertile soils⁷⁷. Other authors from the same region make a distinction between regular vs irregular non-terrace field patterns⁷⁶. A different typology based on social classification of historical field patterns has been recently proposed by Klír²⁴. His extraordinary approach is focused on the social and economic relations of the traditional village society and within the members of this society.

Experts suggest that the diversity of historical field pattern types reflects various factors affecting their genesis, historical development and environmental conditions^{17,39,66}. Several authors have emphasized the influence of age of origin. Žemlička⁸³ claims that there are differences between the field pattern type in ancient settlement area (i.e. regions with settlement continuity since the Neolithic period⁸⁴) and in areas colonized in the medieval period. According to Vermouzek⁸⁵, Klápště⁸⁶, Čulíková⁸⁷ and Šitnerová et al.¹⁵, the irregular, blocky field pattern (segmental plužina; Fig. 1A) is bound to the early medieval period, while narrow and medium-length strips of fields (sectional plužina; Fig. 1C) are characteristic of the high medieval period. Sadravetzová⁸⁸ found segmental plužina predominantly in the high medieval period, and croft plužina (Fig. 1D) in the late medieval and modern period. However, her research was based on dating from written records, which systematically provides dating that is younger than archaeological dating^{89,90}. She also surveyed a sub-mountainous borderland area, which was probably colonized later than the inland of the country. Other authors agree about the influence of age of origin, but without detailed specification (e.g.⁸³). Although the cited references are either small-scale studies^{85,87,88} or review papers/books without a detailed statistical investigation^{15,83,86}, we can conclude that segmental plužina is probably the oldest type, that sectional plužina is younger, and that croft plužina is the youngest type. Examples from Scandinavia show that the time when a regular form of field pattern was introduced varied between regions⁹¹. The transition from segmental plužina to sectional plužina was perhaps caused by the introduction of new agricultural tools, mainly the heavy plough, which required longer parcels^{15,86,92–94}.

Various authors have recognized the existence of a direct relationship between the village layout and the field system type^{15,24,95,96}, but only Sadravetzová⁸⁸ suggests direct links between them (a solitary house or a scattered hamlet—scattered plužina, village green settlements—segmental plužina, a village with hides/*Waldhufendorf* settlement—croft plužina). Many papers and books have presented the expected influence of environmental conditions, mainly terrain properties and soil quality^{15,83,85,97,98} (for further references see⁹⁹), however without providing proper statistical evidence. Lucke et al.¹⁰⁰ report that specific geological conditions (and thus soil properties) affected distinctive field patterns in the Steigerwald region in Germany. Similarly, the occurrence of terraced landscapes in Slovenia has been strongly influenced by geological conditions (mainly flysch and carbonate rocks), by altitude (predominantly 100–600 m), and by southern aspect¹⁰¹; the shape of the Slovenian terraces and their vertical section is also affected by the terrain inclination¹⁰². Unfortunately, these studies did not focus on field pattern types. A clear connection between field pattern type and altitude was found by Sadravetzová⁸⁸ in her local study (segmental and sectional plužina at a lower altitude, croft plužina at higher altitudes). In France, four typical regional forms of terraces have been identified⁶¹.

Based on his earlier micro-regional study¹⁰³, Žemlička⁸³ claims that the number of landlords who simultaneously owned parts of a village in the thirteenth and fourteenth century still affected the field type recorded in the nineteenth century cadastral maps. According to his findings, villages owned by a single landlord in the medieval period mainly still had sectional plužina in the 19th-century maps. In contrast, villages owned by two or more landlords in medieval times had segmental plužina in the nineteenth century. He explains this phenomenon as follows: For the redesign of the field pattern of a village owned by more than one landlord, it was necessary to make a deal among these landlords—which was obviously too complicated. Such villages thus retained their older segmental plužina.

Unfortunately, some of the studies mentioned here are small-scale (local or regional), or lack proper statistical evidence, or do not deal with detailed field pattern typology. However, they suggest possible relationships that can be statistically tested.

The historical field patterns have also been interpreted as a reflection of social structures of the adjacent village society^{24,104,105}. Besides the already discussed environmental factors, Born¹⁰⁶ stresses the influence of agrarian technologies, demography, social and economic factors²⁴. Other authors underline the role of local institutions in shaping historical field patterns^{24,107}.

Spatial distribution of historical field patterns. The spatial extent of historical field patterns has been studied in many European countries: in the Czech Republic^{24,35,108}, France⁶¹, Italy¹⁰⁹, Slovakia^{71,110}, Slovenia^{57,101,111} and historical regions of Saxony (today part of Germany), Silesia (today part of the Czech Republic and Poland), Galicia/Halychyna (today part of Poland and Ukraine)⁶⁵. These studies are on different levels of detail: some sur-




Figure 1. Five types of historical field pattern. (A) Segmental plužina (irregular, blocky structure; Czech: *úseková*), (B) plužina of consolidated/unconsolidated segments (regular, narrow, short strips arranged in many rectangular quadrangles; Czech: *scelené/dělené úseky*), (C) sectional plužina (narrow and medium-length strips parallel to each other, arranged into large irregular blocks; Czech: *traťová*), (D) croft plužina (very long parallel strips following each homestead; Czech: *záhumenicová/dělková*), (E) without internal division, and others. First column: schematic drawing of the field pattern [drawings (A–D) by Šitnerová et al.¹⁵, after Černý⁸¹, drawing E by Václav Fanta, on the base of Černý's⁸¹], second column: *Imperial Imprints of the Stable Cadastre* mid-nineteenth century maps⁸², third column: historic (B,C) or recent (A,D,E) aerial photographs⁸².

veyed the whole area of a country¹¹¹, while others identified only a number of areas with remnants of historical field patterns³⁵. Some authors also tried to determine the spatial distribution of various types of historical field patterns^{24,35,65}, but none succeeded on the level of an entire country.

The focus of our study. In this paper, we decided to focus on the situation in the Czech Republic. This region is characterized by the availability of high-quality historical maps depicting field patterns^{82,112,113}, and also by a long tradition of archaeological research on medieval settlements^{86,114–118}, on landscape and environmental archaeology^{25,28,89,119,120} and on historical field systems^{8,15,28,34,121–126}. The Czech Republic is also characterized by high geological diversity¹²⁷, by diversity of landscape types^{128,129} and by diverse vernacular architecture¹³⁰. Such diversity may have contributed to the historical development of field pattern types.

Within the last millennium, the cultural landscape of Central Europe has undergone several dramatic transformations that have changed the settlement structure: medieval colonization and field pattern redesign^{86,131,132}, the 15th-century religious wars, which destroyed many villages^{133–135}, the Thirty Years' War, which resulted in a decline in population by one third, economic losses, land abandonment and property confiscations^{23,136–140}, eighteenth to nineteenth-century rational redesign of the field ownership structure^{27,85,141}, an industrial revolution with the rapid expansion of cities¹⁴², mid-twentieth century forced collectivization of agriculture, extensive changes in land use, the destruction of traditional field patterns, and ruining of private land ownership by the communist regime^{37,143–148}, as well as land consolidation activities, some of which changed the small-scale historical field pattern into large blocks of arable land¹⁴⁹. At the present time, some parts of the agricultural landscape are being abandoned for various reasons³⁸. Each of these changes has affected the image of the landscape and its functioning.

The plůžina field patterns, which prevailed in the Czech cultural landscape at the time of its medieval colonization and redesign, have gradually changed. Some plůžinas disappeared along with villages that were lost in wars and gradually became re-forested^{125,150}. Others, especially those in the most fertile areas, were transformed into larger arable fields, which can be seen on Stable cadastre maps dating back to the first half of the nineteenth century. However, in the 1950s, plůžina field patterns were still predominant in large parts of the Czech territory⁷⁹. This situation changed fundamentally in the second half of the twentieth century, when many of these landscapes disappeared due to agricultural intensification or due to agricultural extensification and gradual re-forestation².

The rapid loss of plůžinas, along with their outstanding historical, natural and aesthetic values, has led to efforts to gain a deeper understanding of these valuable landscape segments, and to conserve them. However, there still remains a lot that is unknown about plůžinas. As we have noted above, the historical field patterns have different shapes and structuring. Although the role of environmental or cultural/historical factors in determining the type of plůžina prevailing in a specific location has been discussed for decades, precise identification of the “driving forces” supported by statistical evidence is still missing, as is the spatial distribution of individual field pattern types on a nationwide scale. This paper will therefore focus on identifying ecological, geographical and historical factors that could have shaped the spatial structure of the historical field patterns and the spatial distribution of individual types of field patterns within the Czech Republic. In other words, why does the cultural landscape of Central Europe have its present appearance? What formative factors can be statistically linked to its development?

Methods

Data collection. Since it is expected that the field pattern types differed across the centuries^{83,87,88}, we wanted to include the age of the villages as one of the predictors. Because the dating of the origin of historical settlements by widely-available written sources is highly biased^{89,90}, we decided to work with archaeologically dated settlements only. We used a database from our previous research⁸⁹, which contains a subset from the *Archaeological Database of Bohemia*¹⁵¹. This subset includes 527 medieval and early modern age settlements dated by precise archaeological methods (for a detailed description of the data filtering, please refer to the Methods section in the respective paper). In four cases, it was impossible to determine the field pattern types of the respective settlements (either there were no fields in the cadastre, or the old cadastral maps were missing). Such cases were excluded from the dataset. As a result, we worked with 523 settlement sites (for their geographic distribution, see Fig. S1). Unfortunately, archaeological data in sufficient quality and data structure are available only for the western part of the country (the historical region of Bohemia), and not for the entire country. This research therefore covers approximately two-thirds of the Czech Republic only.

The types of field patterns were derived from the *Indication sketches of the Stable Cadastre* (German: *Urmappe des Franziszeischen Katasters*¹⁵², Czech: *indikační skici stabilního katastru*), a set of old cadastral maps from the first half of the nineteenth century⁸². These maps are the oldest historical source available for the whole country depicting the field pattern structure. The scale of these maps is 1 : 2 880, and they are characterized by high geodetical accuracy. These maps, as a highly respected source, have been routinely used in many historical, geographical and ecological studies^{2,28,153–155}.

Within recent decades, experts have identified several typological systems of historical field patterns^{79–81,156,157}. Most typological systems contain six to nine types of field pattern. We took the most distinctive types from Černý's⁸¹ typology, and we grouped other very similar types. As a result, we obtained four basic types (A: segmental, B: consolidated/unconsolidated segments, C: sectional, D: croft plůžina), and we also added one more type (E) for “others/unspecifiable/without internal division” (Fig. 1).

In most cases, the cadastre of a single village contains two or more types of field patterns. In each cadastre, we therefore described all types of field patterns that were present, and then we estimated their relative extent, rounded to tens of percentage points (example: in the village of *Mochov*, the following types of field pattern were identified: type A 50%, type B 30%, type E 20%). Of course, the sum of the relative extents of all types in

one cadastre is 100%. We also recorded the *field pattern diversity* as the number of field pattern types within one cadastre.

All predictors (independent variables) were derived for points representing the centre of the respective village or town. We worked with four environmental variables: *Altitude* data was taken from the SRTM digital terrain model¹⁵⁸. *Terrain undulation* was measured as the average degree of slope within a 4 km radius from the village/town, while the slope was derived from the SRTM model¹⁵⁸. A description of the *geological* bedrock was derived from the geological map¹⁵⁹ and was simplified into four categories (metamorphic / igneous / sedimentary / quaternary). We thank geologist Radana “Radouch” Kavková for her assistance with this simplification. The specific natural *soil fertility* values (0 = worst, 100 = best) were calculated from the soil units data¹⁶⁰, using the methodology described in Bečvářová et al.¹⁶¹.

Next, we operated with historical/geographical variables: The *archaeological dating* was obtained from the Archaeological Database of Bohemia¹⁵¹. The *ancient settlement area* (area inhabited continuously since the Neolithic) was defined as cadasters with Neolithic findings obtained from the Archaeological Database of Bohemia¹⁵¹. We thank archaeologist Jiří “Bumča” Bumerl for his assistance with the archaeological data filtering. To observe the possible impact of important medieval monasteries, we measured the distance to *monasteries* founded in the twelfth and thirteenth centuries^{162,163}. The *Cadastre area* was calculated from the current geographical data¹⁶⁴. *Settlement type* data was derived from the database of historical settlements by Kuča¹⁶⁵. This predictor describes the layout composition of towns and villages in 8 different types (villages with regular hides, orthogonal village green locations, non-orthogonal village green locations, small villages, modern parcel villages and dispersed villages, other villages, modern settlements, historical towns). We grouped the “orthogonal village green locations” and the “non-orthogonal village green locations” into a single group called “village green locations”; we also grouped “small villages”, “modern settlements”, and “other villages” into a single group called “other villages” (for a detailed description of this typology, see¹²⁶). *Confiscates* were derived from Semotanová and Cajthaml¹⁶⁶ as areas with a forced change of landlord in the seventeenth century. [In the first half of the seventeenth century, the emperor from the House of Habsburg confiscated the properties of revolting members of the nobility (primarily protestants) and installed foreign (catholic) nobility in vacant dominions. This process involved a substantial part of the country].

We also added a variable describing historical subsistence strategies: *Agriculture* represents the agricultural regions in the sixteenth century¹⁶⁷, dividing the country into three areas: wheat areas (commercial agriculture production), flax areas (linen production, secondary non-agrarian production), and other areas.

Last but not least, we would like to stress that this list of variables does not include all possible predictors. We think it would be fascinating to have, for example, data on the medieval ownership structure, and to test Žemlička’s⁸³ hypothesis on the influence of the number of landlords. Unfortunately, it is impossible to gather such data for the whole country.

All dependent and independent variables are arranged in Table 1. The geographical data were processed in QGIS 3.22.0 software (<https://qgis.org/en/site/>) and SAGA 7.8.2 software (<https://saga-gis.sourceforge.io/en/index.html>)¹⁶⁸.

Statistical analysis. We assess the effect of our predictors on the dependent variables using regression analyses. The choice of the regression model depends crucially on the nature of the dependent variable. It might feel natural to treat *diversity* (the number of different types of field pattern) as a count outcome; however, its values are restricted to the 1–5 range, due to our classification of field patterns. We therefore adopt two alternative modelling strategies: (i) truncated Poisson regression, with truncation points set at 0 and 6, (ii) an ordinal logistic regression that considers *diversity* a general 1–5 ordinal scale.

In the regressions that explained the relative extent of field patterns A–E, the outcome is a *composition*, or a *multinomial fractional response*. The presence of zeros (in most cadastres, not all patterns are represented) rules out the use of the standard Dirichlet regression approach. We therefore opted for the fractional multinomial logistic quasi-maximum likelihood approach, inspired by Papke and Wooldridge’s¹⁶⁹ estimator for a (single) fractional response and implemented by Buis¹⁷⁰.

Our explanatory variables exhibited a mild degree of collinearity; most notable correlations (0.55–0.62) occurred between environmental predictors (*altitude*, *terrain undulation*, *soil fertility*). These correlations are not strong enough to preclude simultaneous use of the variables in a multivariate regression setting. However, in the fractional multinomial logistic regressions, where four coefficients per predictor are being estimated, the effect of collinearity might be more pronounced; therefore, for robustness, we present the results from both bivariate and multivariate regressions. *Geology* was excluded from the multivariate regressions due to a nearly 20% rate of missingness; for the other variables, values were missing in less than 2% of the sample (see Table 2 for sample sizes in bivariate regressions). Several numeric predictors (*terrain undulation*, *altitude*, *cadastre size*, *monasteries*) showed a substantial positive skew; we log-transformed their values prior to all statistical analyses.

Finally, we used the *relative variable importance* (RVI) measure to rank predictors in terms of the weight of evidence for their inclusion in the best model. A predictor’s RVI, also referred to as the parameter weight, the parameter inclusion probability or the proportion of evidence^{171–173}, is obtained by (i) running the regressions with all possible subsets of the available predictors, (ii) calculating the Akaike weight for each estimated model, and (iii) summing the Akaike weights across all models that included the predictor of interest. RVI is bounded between 0 and 1; thresholds of 0.4¹⁷¹ or 0.5¹⁷⁴ have been recommended to separate important variables and unimportant variables.

All regression analyses were conducted in Stata 17.0 (<https://www.stata.com/>). The user-contributed commands *fmlogit*¹⁷⁰, *miinc*¹⁷³, and *estout*¹⁷⁵ were used for the fractional multinomial logistic regression, for the RVI

Variable name	Description	Data source
Dependent variables (response)		
<i>Diversity</i>	Number of field pattern types within one cadastre	Land Survey Office ⁸²
<i>Type A</i>	The relative extent of the respective field pattern type in the cadastre (0–100 %)	
<i>Type B</i>		
<i>Type C</i>		
<i>Type D</i>		
<i>Type E</i>		
Environmental independent variables (predictors)		
<i>Altitude</i>	Meters above sea level	GISAT ¹⁵⁸
<i>Terrain undulation</i>	Average degree of slope within a 4 km radius	
<i>Geology</i>	Geological bedrock, simplified	Czech Geological Survey ¹⁵⁹
<i>Soil fertility</i>	Relative soil fertility (0 = worst, 100 = best)	State Land Office ¹⁶⁰ , Bečvářová et al. ¹⁶¹
Historical/geographical independent variables (predictors)		
<i>Archaeological dating</i>	Age of the settlement	Institute of Archaeology of the Czech Academy of Sciences Prague ¹⁵¹
<i>Ancient settlement area</i>	Presence or absence of Neolithic findings in the cadastre	Institute of Archaeology of the Czech Academy of Sciences Prague ¹⁵¹
<i>Monasteries</i>	Distance to medieval monasteries	Purš ¹⁶² , Hrnčiarová et al. ¹⁶³
<i>Confiscates</i>	Areas confiscated in the seventeenth century	Semotanová and Cajtham ¹⁶⁶
<i>Settlement type</i>	Type of layout composition	Kuča ¹⁶⁵
<i>Cadastre area</i>	Square meters	Arcdata Praha ¹⁶⁴
Historical subsistence strategies		
<i>Agriculture</i>	agricultural/subsistence strategies in sixteenth century	Klír ¹⁶⁷

Table 1. All variables used in the study.

Dependent variable	Proportions of field patterns			Field pattern diversity			
	Fractional multinomial logit			Ordinal logit		Truncated Poisson	
	Model	<i>N</i>	<i>p</i> -value	RVI	<i>p</i> -value	RVI	<i>p</i> -value
Terrain undulation (log)	523	< 0.0001	0.998	0.509	0.284	0.025	0.280
Altitude (log)	523	0.0310	0.282	0.116	0.609	0.016	0.423
Soil fertility	516	0.0110	0.014	0.594	0.337	0.023	0.303
Cadastre area (log)	523	< 0.0001	0.588	< 0.001	0.997	< 0.001	0.950
Archaeological dating	523	< 0.0001	0.050	0.046	0.589	0.010	0.393
Monasteries (log)	523	0.1352	0.034	0.708	0.290	0.026	0.274
Confiscates	515	0.0013	0.212	0.638	0.267	0.024	0.284
Settlement type	514	< 0.0001	1.000	0.041	0.064	0.019	0.054
Geology	405	< 0.0001		0.661		0.146	

Table 2. Summary statistics for bivariate and multivariate regressions. *N* the number of observations in bivariate regressions; *p*-value overall model *p*-value in bivariate regressions; *RVI* relative variable importance, i.e., sum of the Akaike weights across all models containing the given covariate in the all-subset regressions (*geology* was not included in the all-subset regressions).

calculation, and for the regression tables, respectively. The geographical interpolation for Fig. 4 was performed in SAGA software¹⁷⁶ using the *universal kriging* tool.

Results

According to Table 2, the most significant variables affecting the proportions of field pattern types were *settlement type* (*RVI* = 1.000), *terrain undulation* (*RVI* = 0.998), and *cadastre area* (*RVI* = 0.588). These variables affected almost all field pattern types. The influence of other variables had less impact on the overall variability. Some variables did not affect any field pattern type at all: distance to medieval *monasteries* (see Table 2) and *ancient settlement area* (*p*-value from the bivariate regression is 0.092). The field pattern diversity was influenced by one variable only: the *cadastre area* (*RVI* = 0.950).

Field pattern *type A* (segmental *plužina*) was associated with very rugged *terrain* ($p < 0.001$), small *cadastre area* ($p < 0.001$), preferably igneous but definitely non-sedimentary *geological* bedrock (see Fig. 3) and specific

types of *settlement layout type* (“village green locations” and “other”). The influence of *altitude* was insignificant; however, Fig. 2 suggests a possible preference for higher locations. Both Table 3 and Fig. 2 agree with the preference for poor *soil fertility* of this type, but the *p*-value is above 0.05. According to Fig. 2, this type was connected either with very old (pre-1200 AD) or with very young (post-1500 AD) *archaeological dating*. It was not affected by *confiscates*.

Field pattern *type B* (plužina of consolidated/unconsolidated segments) was associated with flat *terrain*, and with specific types of *settlement layout type* (“village green locations”, “towns” and “other”), and with dominions that remained in possession of the original landlords after the Thirty Years’ War (variable *confiscates*). The *soil fertility* might play a role (a slight preference for more fertile soils). This type was not affected by *altitude* (the proportion of type B within the cadasters seems to be independent of the altitude), *geology*, *archaeological dating* (the percentage of type B did not change over the centuries) or *cadastre area* (clearly insignificant).

Field pattern *type C* (sectional plužina) was associated with flat *terrain* ($p < 0.001$), big *cadastre area*, dominions *confiscated* in the seventeenth century, older (mostly pre 1300 AD) *archaeological dating*, good *soil fertility*, specific types of *settlement layout type* (“village green locations”, “towns” and “other”), and preferably metamorphic *geological bedrock*. The influence of *altitude* is unclear: in Table 3, the bivariate model shows insignificant negative dependence, while the multivariate model shows significant positive dependence, and Fig. 2 suggests relatively mild negative dependence.

Field pattern *type D* (croft plužina) was primarily associated with one specific type of *settlement layout type* (“villages with hides”) and with sedimentary *geological bedrock*. The other variables are problematic, since the bivariate and multivariate models offer different results (see Table 2). Nevertheless, based on Fig. 2, we can say that there was probably a preference for higher *altitude* (the graph suggests that areas over 300–500 m have a higher percentage of type D than lower areas), more *undulating terrain*, poor *soil fertility* (according to the bivariate model, this was apparently quite an important variable), very young (late) *archaeological dating* (post-1300–1400 AD), and maybe the dominions *confiscated* in the seventeenth century. This type was not affected by the *cadastre size*.

Field pattern *type E* (without internal division, and others) was associated with lower *altitude*, igneous *geological bedrock* and specific types of *settlement layout* (all except “villages with hides”). This field pattern was not affected by *terrain undulation*, *soil fertility*, *archaeological dating*, *confiscates* or *cadastre size*.

According to Table 4, the *diversity* was positively affected by the *cadastre area* (a more extensive cadastre area led to higher field pattern diversity). Fig. S2 suggest a possible but insignificant negative influence of *archaeological dating* and *altitude* (older sites and lower altitude should lead to greater diversity). The other variables had no significant effect.

Discussion

The results show that the number of field pattern types within a single cadastre (variable *diversity*) was significantly affected by one predictor only—the size of the cadastre, i.e. the bigger the cadastre area, the greater the diversity. The explanation for this phenomenon is trivial: given the approximately constant count of field pattern types per unit area, a bigger cadastre area offers a higher probability of the occurrence of a specific field pattern type—cf. the “species-area relationship” in ecology¹⁷⁷, which is however driven by other processes. Surprisingly, no other predictors had a significant effect. Nevertheless, we have identified a remarkable spatial distribution of field pattern diversity across the country (Fig. 4F). Some regions are characterized by relatively high diversity (marked by red colour), while others are curiously homogenous (blue). We hypothesize that this distribution may be affected by other factors not included in our study. We can summarize that the average number of field pattern types within a cadastre varies between two and three. During the pre-industrial period, the cadastre size also directly affected the number of inhabitants in a village by providing the area for agricultural activities, i.e. for their subsistence¹³⁷.

Field pattern *type A* (segmental plužina) was characterized by extreme environmental conditions (higher altitude, extremely rugged terrain, igneous bedrock and low soil fertility). Type A is also associated with tiny cadastres and specific types of settlement layouts (“other”)—however, the latter association is insignificant. Field pattern *type A* is situated at settlements founded either in the early medieval period (pre-1200 AD) or in the modern period (post-1500 AD). It is present primarily in a belt across the whole country (Fig. 4A), which is (in most cases) characterized by harsh conditions. We can summarize that type A (segmental plužina) is present in locations with extreme conditions—environmental and cultural (cadastre size). It is a kind of “extremophile organism”—it can “survive” in situations where other strategies wither away^{178–180}. We hypothesize that this field pattern type could initially also have been present in areas with better conditions, but it may have been replaced by type C (sectional plužina), because of type C’s better suitability for medieval agriculture^{132,181}. Type A then remained only in areas with bad conditions for agriculture. Type A was frequent at settlements founded in the early medieval period, which corresponds well with previous findings^{15,85–88}. The second peak of this type lies in the early modern period. We think that this reflects the late medieval to early modern colonization of highlands and sub-mountainous regions, where other types of field patterns could not be established because of the harsh environmental conditions and limited spatial possibilities. Surprisingly, no terraces (as we know them from the Mediterranean area) have been built (except for the vineyards in Moravia) in the areas with significant terrain undulation (up to 10°). Perhaps it was easier to organize segmental plužina structure than to build terrace walls.

Field pattern *type B* (plužina of consolidated/unconsolidated segments) was characterized by relatively high soil fertility, flat terrain and specific types of settlement layout (mostly “towns”). This type occurs at almost all altitudes and in almost all ages within the study period. The geology and the cadastre area were also insignificant. It is present primarily in southern Bohemia (Fig. 4B). To conclude, this field pattern type prefers relatively flat terrain and higher soil fertility, i.e. good conditions for agriculture. Other variables have limited influence. The




Figure 2. Influence of continuous variables. The percentages refer to the proportion of the specific field pattern type in the examined cadastres. The descriptions of variables and their units of the x-axes are on the left side of the respective panels.

Regression	A: segmental		B: (un)cons. segments		C: sectional		D: croft		E: others	
	AME	SE	AME	SE	AME	SE	AME	SE	AME	SE
Terrain undulation (log)										
Bivariate	11.51***	(0.0239)	-8.253**	(0.0260)	-9.605***	(0.0212)	6.503**	(0.0214)	-0.158	(0.0196)
Multivariate	14.36***	(0.0305)	-10.33**	(0.0353)	-11.39***	(0.0306)	2.922	(0.0235)	4.445	(0.0253)
Altitude (log)										
Bivariate	4.320	(0.0383)	-2.210	(0.0424)	-1.873	(0.0347)	7.029*	(0.0289)	-7.266*	(0.0326)
Multivariate	-9.482	(0.0487)	9.951	(0.0591)	13.55**	(0.0521)	-3.768	(0.0388)	-10.25*	(0.0419)
Soil fertility										
Bivariate	-13.29	(0.0790)	13.00	(0.0892)	16.44*	(0.0724)	-20.64*	(0.0802)	4.490	(0.0585)
Multivariate	-10.42	(0.110)	0.845	(0.125)	11.86	(0.106)	0.0803	(0.0887)	-2.368	(0.0796)
Cadastre area (log)										
Bivariate	-7.339***	(0.0185)	-1.146	(0.0197)	4.896**	(0.0165)	3.625	(0.0192)	-0.0358	(0.0120)
Multivariate	-6.139***	(0.0182)	-3.239	(0.0214)	6.209***	(0.0188)	0.814	(0.0207)	2.354	(0.0157)
Archaeological dating										
Bivariate	1.287	(0.0114)	-0.0455	(0.0106)	-3.123***	(0.00857)	2.980***	(0.00824)	-1.097	(0.00725)
Multivariate	0.656	(0.0120)	0.817	(0.0122)	-1.507	(0.0101)	1.027	(0.00803)	-0.993	(0.00759)
Monasteries (log)										
Bivariate	-1.612	(0.0154)	-0.491	(0.0176)	-2.628	(0.0143)	3.677	(0.0201)	1.054	(0.0143)
Multivariate	-1.413	(0.0154)	-0.0109	(0.0203)	-1.734	(0.0163)	1.945	(0.0146)	1.212	(0.0136)
Confiscates										
Bivariate	-3.531	(0.0269)	-6.280*	(0.0310)	5.533*	(0.0264)	7.376**	(0.0239)	-3.097	(0.0204)
Multivariate	-1.215	(0.0265)	-6.314*	(0.0317)	5.939*	(0.0258)	3.827	(0.0201)	-2.237	(0.0206)
Settlement type										
<i>Village with hides</i>	Ref.		Ref.		Ref.		Ref.		Ref.	
<i>Village green</i>										
Bivariate	9.002*	(0.0434)	15.45**	(0.0555)	21.01***	(0.0509)	-53.19***	(0.0542)	7.720**	(0.0285)
Multivariate	10.39*	(0.0460)	10.87	(0.0591)	18.06***	(0.0525)	-46.77***	(0.0571)	7.453*	(0.0316)
<i>Dispersed</i>										
Bivariate	15.08	(0.0791)	14.72	(0.0781)	1.032	(0.0554)	-43.12***	(0.0782)	12.28*	(0.0571)
Multivariate	10.99	(0.0747)	10.34	(0.0775)	4.493	(0.0658)	-39.68***	(0.0760)	13.86*	(0.0629)
<i>Towns</i>										
Bivariate	5.730	(0.0386)	22.54***	(0.0485)	14.44***	(0.0430)	-49.43***	(0.0539)	6.722**	(0.0241)
Multivariate	7.993*	(0.0398)	23.89***	(0.0553)	8.672*	(0.0417)	-44.95***	(0.0547)	4.392	(0.0271)
<i>Others</i>										
Bivariate	15.45***	(0.0353)	13.26**	(0.0406)	9.238*	(0.0364)	-52.71***	(0.0499)	14.76***	(0.0238)
Multivariate	13.14***	(0.0347)	10.51*	(0.0470)	10.39**	(0.0394)	-47.04***	(0.0550)	12.99***	(0.0275)
Geology										
<i>ign</i>	Ref.		Ref.		Ref.		Ref.		Ref.	
<i>met</i>										
Bivariate	-9.048	(0.0536)	1.019	(0.0525)	10.74**	(0.0414)	1.238	(0.0307)	-3.945	(0.0384)
<i>qua</i>										
Bivariate	-8.944	(0.0601)	3.976	(0.0625)	4.873	(0.0453)	2.946	(0.0423)	-2.851	(0.0467)
<i>sed</i>										
Bivariate	-14.63**	(0.0510)	-2.842	(0.0510)	5.688	(0.0387)	20.52***	(0.0399)	-8.739*	(0.0365)

Table 3. Average marginal effects from bivariate and multivariate regressions: fractional multinomial logit explaining the proportions of field patterns. (i) To enhance readability, *archaeological date* and *soil fertility* were divided by 100 prior to running the regressions. (ii) the standard errors, in parentheses, were obtained from regression estimates via the delta method. (iii) * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

shape of this field pattern testifies to rational land consolidation in restricted blocks without significant spatial requirements or splitting the larger fields, initially owned by the nobility, into small fields during the modern period^{27,182,183}. As field pattern *type B* is not bound to a specific historical period, it was probably a general method of secondary land consolidation in areas with good agricultural conditions^{182,184}.

Field pattern *type C* (sectional plužina) was characterized by optimal conditions for agriculture (lower and medium altitude, flat terrain, very high soil fertility, metamorphic bedrock, and big cadastre area). It is associated with settlements established in the early and high medieval period (pre-1300 AD, almost no occurrence after 1400

Regression	Ordinal logit		Truncated Poisson	
	Coeff.	SE	Coeff.	SE
Terrain undulation (log)				
Bivariate	-0.0963	(0.146)	-0.0254	(0.0688)
Multivariate	0.0657	(0.739)	0.0256	(0.780)
Altitude (log)				
Bivariate	-0.364	(0.232)	-0.103	(0.109)
Multivariate	-0.438	(0.182)	-0.152	(0.322)
Soil fertility				
Bivariate	0.252	(0.472)	0.0347	(0.219)
Multivariate	0.101	(0.882)	-0.0139	(0.964)
Cadastral area (log)				
Bivariate	0.450***	(0.110)	0.162**	(0.0507)
Multivariate	0.401**	(0.002)	0.134*	(0.026)
Archaeological dating				
Bivariate	-0.116*	(0.0580)	-0.0345	(0.0267)
Multivariate	-0.101	(0.134)	-0.0259	(0.399)
Monasteries (log)				
Bivariate	0.0373	(0.0996)	0.00564	(0.0467)
Multivariate	0.0611	(0.566)	0.0184	(0.714)
Confiscates				
Bivariate	0.0781	(0.166)	0.0571	(0.0766)
Multivariate	0.0281	(0.871)	0.0333	(0.678)
Settlement type				
<i>Village with hides</i>	Ref.		Ref.	
<i>Village green</i>				
Bivariate	0.164	(0.333)	0.0983	(0.160)
Multivariate	0.138	(0.694)	0.103	(0.536)
<i>Dispersed</i>				
Bivariate	0.146	(0.461)	0.0738	(0.234)
Multivariate	0.398	(0.409)	0.154	(0.525)
<i>Towns</i>				
Bivariate	0.745*	(0.309)	0.303*	(0.148)
Multivariate	0.441	(0.172)	0.202	(0.189)
<i>Others</i>				
Bivariate	0.138	(0.278)	0.0822	(0.138)
Multivariate	0.178	(0.548)	0.112	(0.437)
Geology				
<i>ign</i>	Ref.		Ref.	
<i>met</i>				
Bivariate	0.350	(0.285)	0.0964	(0.132)
<i>qua</i>				
Bivariate	0.166	(0.330)	0.0291	(0.153)
<i>sed</i>				
Bivariate	0.211	(0.281)	0.0401	(0.130)

Table 4. Coefficient estimates for bivariate and multivariate regressions explaining field pattern diversity. (i) To enhance readability, *archaeological date* and *soil fertility* were divided by 100 prior to running the regressions. (ii) * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

AD) and with specific types of settlement layouts (“village green locations”). It is present primarily in western Bohemia (Fig. 4C). This field pattern has much better environmental conditions for agricultural production than type B. It also requires much more space—usually the whole area of the cadastral. Considering the relatively early dating (pre-1300 AD), we suggest two possible ways in which this field pattern was set up: either it was set up in the early medieval period (together with the foundation of the village—see^{15,86,87}), or, like type B, it may have been an instrument of secondary land consolidation^{132,182,185,186}, motivated by excellent conditions for agriculture.

Field pattern *type D* (croft pluzina) was characterized by relatively poor conditions for agriculture (higher altitude, more undulating terrain, sedimentary bedrock, and poor soil fertility). This type is found in settlements established primarily in the late medieval and modern periods (post-1300–1400 AD), corresponding

with Sadravetzová⁸⁸. Type D is present almost exclusively under these particular conditions. It is present mainly in the mountains of northern and eastern Bohemia (Fig. 4D). This type is also strictly bound to “villages with regular hides” and to areas with secondary non-agrarian production (flax areas, Fig. 3). The sedimentary bedrock suggests that it occurred primarily in valleys. We can conclude that this was the last type of field pattern to be set up, as a result of the colonization of primarily mountainous regions. Like segmental plužina, type D offers a more recent adaptation to harsh environmental conditions. As the abundance of this type was strictly driven by the village layout, we assume that it was, in most cases, a result of planned colonization, without later land consolidation.

Field pattern *type E* (without internal division, and others) was characterized by a slightly lower altitude, while no other factors showed a significant effect. This type is present almost throughout the study area (with the exception of southern Bohemia, where there is a high occurrence of type B; Fig. 4E). Based on the preferred environmental conditions, we can describe this type as a “generalist”—it can tolerate a wide range of conditions without specific requirements. This is in accordance with the ecological observation that generalists usually do not need resource-rich habitats¹⁸⁷. This type probably represents land owned by the nobility^{24,182}.

Our findings show an evident link between the selected field pattern type and the settlement layout type (Table 3, Fig. 3). There is an apparent relationship between “villages with regular hides” and croft plužinas (notably, croft plužinas occur only in “villages with regular hides”). “Village green locations” are mostly connected with sectional plužina. “Towns” are primarily (although insignificantly) associated with plužina of (un)consolidated segments. “Dispersed” and “other” are not field-pattern-type specific. These findings partially correspond with previous research by Sadravetzová⁸⁸: we have found an exact link between “villages with regular hides” and croft plužina. Unlike Sadravetzová, whose paper linked “village green locations” to segmental plužina, our results link “village green locations” to sectional plužina. This discrepancy may be caused by the different study areas (local vs. national). The linkage between settlement layout type and field pattern type was also supported by Dohnal and Škabrada¹⁸⁸, who identified a clear relationship between the courtyard size and its hide width. Based on the sequence of houses and field parcels, similar results have also been reported from Northern Europe^{91,189–191}.

Our data suggest a remarkable result (Table 2, Fig. 3): areas confiscated after the Thirty Years’ War have significantly different field pattern types (type C and maybe also type D) than areas that remained in the possession of their original landlords (type B). Types B and C occur in similar environmental conditions (high soil fertility, flat terrain). These two types could be different adaptations to the same requirements: rational, orthogonal organization of parcels. The difference between them lies in their spatial extent. While type B (plužina of (un)consolidated segments) requires only a limited area (e.g. 5% of the whole cadastre), type C (sectional plužina) needs at least one half of the cadastre area. We propose the following explanation of the observed data: As the confiscated cadasters were sold to new (foreign and wealthy) nobility, the new landlords probably had much more financial resources, foreign experience and willingness to reorganize the field pattern than the original owners [example: activities of Count von Sporck, son of the emperor’s general and a member of the “new” nobility, on his estates in Central and Eastern Bohemia^{192,193}]. The new nobility was also catholic (as was the emperor), and the (primarily protestant) peasants may have been afraid to revolt in the times when the counter-reformation was beginning. These results also indicate that land consolidation activities also took place in the modern period, probably as a result of various land reforms¹⁴¹.

Contrary to the expectations suggested by Žemlička⁸³, the variable *ancient settlement area* showed no significant effect. In other words, the presence of continuous settlement activities since the Neolithic played no role. This finding may seem rather surprising, as it has been considered that ancient settlement area had shared characteristics¹²⁹.

Similarly, distance to *monasteries* was also an insignificant variable. Nevertheless, we cannot completely exclude the possible influence of medieval monasteries, which had some of their estates in distant parts of the country¹⁹⁴.

Interestingly, several conditions are characteristic of one type of field pattern only. E.g. “villages with regular hides” are (in most cases) surrounded by croft plužinas only; extremely high terrain undulation is a precise predictor for segmental plužina; huge cadastres indicate the occurrence of sectional plužina, etc. Likewise, several regions are characterized by only one type of field pattern (Fig. 4). This finding is similar to observations from France, where typical regional forms of terraced landscapes have been identified⁶¹. Maybe the field pattern type was part of the specific regional traditions (like the vernacular architecture and the traditional folk costumes—cf.¹⁹⁵)?

Our findings on the environmental predictors of type A (segmental plužina) are in contradiction to findings presented by Klír²⁴ in his local study from western Bohemia. According to his results, the segmental plužina prefers the optimal conditions for agriculture. We think this discrepancy might be caused by local history or specific conditions in the region.

We should mention here several possible limitations of our study. First, the settlement selection was not random, but was based on the availability of archaeological data (see Fig. S1). This could theoretically pose a bias towards selecting specific geographical areas, and could thus affect the final results. Second, the field pattern types were determined by human researchers, and humans can make mistakes. Although we checked our data properly, we cannot exclude the possibility that the type of field pattern may have been categorized incorrectly in some cases. However, this should not play a role due to the relatively big sample. Third, the spatial extent of our study was limited to the western part of the Czech Republic by the availability of archaeological data. The generalization of our results is therefore disputable. It would open quite new possibilities to compare our results with findings from other countries, which is nevertheless dependent on the availability of standardized datasets.




Figure 3. Influence of categorical variables. The percentages refer to the proportion of the specific field pattern type in the examined cadastres. The descriptions of variables of the x-axes are on the left side of the respective panels.




Figure 4. (A–E) Geographical distribution of field pattern types within Bohemia (western part of the Czech Republic). The percentages refer to the proportion of each field pattern type in the examined cadastres. The field pattern types are shown in the schematic black-and-white drawings. (F) Diversity of field pattern types, i.e. the number of field pattern types within examined cadastres. All panels: The original point-based data were interpolated using the *universal kriging* interpolation tool in SAGA software¹⁷⁶. The eastern part of the Czech Republic (Moravia, Silesia) is not represented because of missing archaeological data. The black-and-white schematic drawings of plužina types in panels (A–D) are from Šitnerová et al.¹⁵, after Černý⁸¹. The map was created by authors using QGIS software (<https://qgis.org/en/site/>, version 3.22.0) and SAGA software (<https://saga-gis.sourceforge.io/en/index.html>, version 7.8.2).

Conclusion

Our findings confirm the influence of environmental conditions, mostly terrain undulation and soil fertility, and the effect of geographical/cultural variables (e.g. cadastre size) on the field pattern type prevailing in a specific location. Moreover, we can suggest various “ecological adaptations” to specific environmental conditions (cf.¹⁹⁶): Sectional plužina and plužina of (un)consolidated segments prefer good conditions for agriculture. Segmental and croft plužinas can withstand even very harsh conditions, and type E (others) is not linked to the conditions important for agriculture. These adaptations differ in the spatial extent of the field pattern type within a cadastre (sectional plužina vs. plužina of (un)consolidated segments); or age of origin, regional traditions and customary settlement layout type (segmental plužina vs. croft plužina) may have also contributed to the observed distribution. As suggested by previous studies, there is a clear link between field pattern type and settlement layout type. However, the link is valid for limited cases only (“village with regular hides”—croft plužina, “village green locations”—sectional plužina). This corresponds with ethnographic findings^{91,188}. On the other hand, some older hypotheses (e.g. the influence of an ancient settlement area, proposed by Žemlička⁸³) were not supported by our data.

Besides affecting the size of the village by providing an area for agriculture¹³⁷, the size of the cadastre is also responsible for the number of field pattern types within the cadastre—and is therefore linked with cultural landscape diversity. The settlement density (i.e. the distance between villages) thus affected the village size and the overall landscape structure.

The type of field pattern has also been affected by seventeenth century confiscations: cadasters in confiscated estates have significantly different field pattern types than cadasters which remained in the possession of the original landlords. This phenomenon may have been caused by the non-local economic knowledge and the access to resources of the new nobility, allowing for re-organization of older field systems in accordance to the preferences of the impending landowners.

The spatial distribution of field pattern types within the country was probably also driven by other factors that were not included in our study. Strong regional prevalences of specific field pattern types can be observed (Fig. 4), which cannot be explained by environmental or geographical variables only. Theoretically, we could hypothesize that the field pattern type was a part of specific local/regional traditions (cf.⁶¹)—and that the adaptation to similar environmental (and other) requirements could differ from region to region.

Our findings do not dispute the results of previous historical research. Instead, we have confirmed earlier findings and have put them into a new, broader context. Together with historians and archaeologists, we can summarize that: During the early medieval period, the earliest plužina types were probably the segmental types (type A). Some plužinas were re-designed to sectional types (type C) during the high-medieval colonization (especially in areas with good conditions for agriculture). During the whole studied period (i.e. in medieval

and early modern times), pluzinas of consolidated/unconsolidated segments (type B) were popular as a rational way to consolidate the land-ownership structure. Croft pluzinas (type D) were used during the late medieval colonization of sub-mountainous areas.

Apart from the environmental factors, our study shows a significant influence of historical/social factors (cadastre size, seventeenth century confiscations, settlement layout type, regional clustering). These predictors point to various historical turbulences through the centuries and their impact on the cultural landscape at the detailed scale.

To the best of our knowledge, our paper provides the first detailed analysis of the geographical distribution of traditional field systems at the scale of an entire state (the historical Kingdom of Bohemia). In addition to the environmental predictors, we want to stress the importance of social factors—e.g. the relationship with the settlement layout type, with the cadastre size, with the war disturbances and with the subsequent social unrest in the seventeenth century. It is crucial to study both types of driving forces and to take into account the mutual impacts of environmental and social factors.

Data availability

All used data are available in Dataset S1.

Received: 19 January 2022; Accepted: 9 May 2022

Published online: 23 May 2022

References

1. UNESCO. Operational Guidelines for the Implementation of the World Heritage Convention. *World Heritage Centre* <https://whc.unesco.org/document/178167> (2019).
2. Sklenicka, P. et al. Remnants of medieval field patterns in the Czech Republic: Analysis of driving forces behind their disappearance with special attention to the role of hedgerows. *Agric. Ecosyst. Environ.* **129**, 465–473 (2009).
3. Rippon, S. *Historic Landscape Analysis: Deciphering the Countryside* (Council for British Archaeology, 2008).
4. *Historické kulturní krajiny České republiky [Historical cultural landscapes of the Czech Republic]*. (Výzkumný ústav Silva Taroucy pro krajinu a okrasné zahradnictví, v. v. i., 2020).
5. Meier, D. Man and environment in the marsh area of Schleswig-Holstein from Roman until late Medieval times. *Quat. Int.* **112**, 55–69 (2004).
6. Baudry, J., Bunce, R. G. H. & Burel, F. Hedgerows: An international perspective on their origin, function and management. *J. Environ. Manage.* **60**, 7–22 (2000).
7. Burel, F. & Baudry, J. Social, aesthetic and ecological aspects of hedgerows in rural landscapes as a framework for greenways. *Landsc. Urban Plan.* **33**, 327–340 (1995).
8. Bayer, T. & Beneš, J. Středověká terasová pole na Šumavě jako hydropedologický fenomén a archeologický problém. *Archeol. rozhl.* **56**, 139–159 (2004).
9. Merot, P. The influence of hedgerow systems on the hydrology of agricultural catchments in a temperate climate. *Agronomie* **19**, 655–669 (1999).
10. Ucar, T. & Hall, F. R. Windbreaks as a pesticide drift mitigation strategy: A review. *Pest Manag. Sci. Former. Pestic. Sci.* **57**, 663–675 (2001).
11. Marshall, E. J. P. & Moonen, A. C. Field margins in northern Europe: Their functions and interactions with agriculture. *Agric. Ecosyst. Environ.* **89**, 5–21 (2002).
12. Caubel-Forget, V., Grimaldi, C. & Rouault, F. Contrasted dynamics of nitrate and chloride in groundwater submitted to the influence of a hedge. *Comptes Rendus l'Académie des Sci. Ser. IIA Earth Planet. Sci.* **332**, 107–113 (2001).
13. Forman, R. T. T. & Godron, M. *Landscape Ecology* (Wiley, 1986).
14. Deckers, B., Kerselaers, E., Gulincx, H., Muys, B. & Hermy, M. Long-term spatio-temporal dynamics of a hedgerow network landscape in Flanders, Belgium. *Environ. Conserv.* **32**, 20–29 (2005).
15. Šitnerová, I. et al. Archeologický výzkum pluzin a zemědělských teras jako fenoménu historické krajiny České republiky [Archaeological research into field systems and agrarian terraces as the phenomena of the historical landscape in the Czech Republic]. *Archaeol. hist.* **45**, 141–165 (2020).
16. Beneš, J. & Zvelebil, M. A historical interactive landscape in the heart of Europe: The case of Bohemia. In *The archaeology and Anthropology of Landscape Shaping your Landscape* (eds Ucko, P. J. & Layton, R.) 73–93 (Routledge, 1999).
17. Aston, M. *Interpreting the Landscape. Landscape Archaeology and Local History* (Routledge, 1985).
18. Collier, M. J. Are field boundary hedgerows the earliest example of a nature-based solution?. *Environ. Sci. Policy* **120**, 73–80 (2021).
19. Deutscher, J. The visual beauty of linear green-structures in the rural landscape, principles of aesthetic evaluation and its application in landscape management. in *Public recreation and landscape protection: with man hand in hand?* (eds Fialova, J. & Pernicova, D.) (2014).
20. Sklenicka, P., Kottová, B. & Šálek, M. Success in preserving historic rural landscapes under various policy measures: Incentives, restrictions or planning?. *Environ. Sci. Policy* **75**, 1–9 (2017).
21. Sklenička, P. & Pittnerová, B. Pozemkové úpravy v územích s dochovanou středověkou pluzinou. *Pozem. úpravy* **51**, 19–20 (2005).
22. *Povodně a sucho. Krajina jako základ řešení. Sborník příspěvků ze seminářů Komise pro životní prostředí Akademie věd ČR konaných ve dnech 8. října 2013 a 5. června 2014.* (Botanický ústav Akademie věd České republiky, v. v. i., 2014).
23. Poschod, P. *Geschichte der Kulturlandschaft [History of the Cultural Landscape]*. (Ulmer, 2015).
24. Klír, T. *Rolnictvo na pozdně středověkém Chebsku. Sociální mobilita, migrace a procesy pustnutí.* (Univerzita Karlova, 2020).
25. Gojda, M. *Archeologie krajiny. Vývoj archetypů kulturní krajiny [Archaeology of Landscape. The Evolution of Cultural Landscape Archetypes]*. (Academia, 2000).
26. Klír, T. Rekonstrukční mapy polních systémů [Reconstruction maps of field systems]. *Hist. geogr.* **34**, 261–295 (2007).
27. Dohnal, M. *Historická kulturní krajina v novověku. Vývoj vsi a pluziny v Borovanech u Bechyně [A historic cultural Landscape in the early modern period. Settlement and field system evolution at Borovany u Bechyně]*. (Ústav archeologické památkové péče středních Čech, 2003).
28. Houfková, P. et al. Origin and development of long-strip field patterns: A case study of an abandoned medieval village in the Czech Republic. *CATENA* **135**, 83–91 (2015).
29. Pešta, J. Několik poznámek ke studiu půdorysné struktury venkovských sídel na území Čech: Einige Bemerkungen zum Studium der Grundriss-Struktur der ländlichen Sitze auf dem Gebiet Böhmens. *Průzkumy památek* **7**, 153–168 (2000).
30. Roberts, B. K. *Landscape of Settlement. Prehistory to Present* (Routledge, 1996).

31. Chang, P. Archaeological landscapes: The ethnoarchaeology of pastoral land use in the Grevena province of northern Greece. In *Space, Time, and Archaeological Landscapes* (eds Rossignol, J. & Wandsnider, L.) 65–89 (Springer, 1992).
32. Gadot, Y. *et al.* OSL dating of pre-terraced and terraced landscape: Land transformation in Jerusalem's rural hinterland. *J. Archaeol. Sci. Reports* **21**, 575–583 (2018).
33. Turner, S. *et al.* Agricultural terraces in the Mediterranean: Medieval intensification revealed by OSL profiling and dating. *Antiquity* **95**, 773–790 (2021).
34. Štínerová, I. *et al.* Landscape transformed: Archaeological, historical and environmental dating of the early modern field system in Valštejn, Czech Republic. *Interdiscip. Archaeol. Nat. Sci. Archaeol.* **11**, 089–101 (2020).
35. Kuča, K. Oblasti dochovaných strukturálně výrazných plujin v České republice [Areas of extant significant historical field patterns in the Czech Republic]. *Zprávy památkové péče* **74**, 34–49 (2014).
36. Kremenič, T., Andlar, G. & Varotto, M. How did sheep save the day? The role of dry stone wall heritage and agropastorality in historical landscape preservation. A case-study of the town of cres olive grove. *Land* **10**, (2021).
37. Lieskovský, J. *et al.* Factors affecting the persistence of traditional agricultural landscapes in Slovakia during the collectivization of agriculture. *Landsc. Ecol.* **29**, 867–877 (2014).
38. Lieskovský, J. *et al.* The abandonment of traditional agricultural landscape in Slovakia: Analysis of extent and driving forces. *J. Rural Stud.* **37**, 75–84 (2015).
39. Lebeau, R. *Les grands types de structures agraires dans le monde.* (Masson – Armand Colin, 1996).
40. Pregill, P. & Volkman, N. *Landscape in History. Design and Planning in the Eastern and Western Traditions* (Wiley, 1999).
41. *World Terraced Landscapes: History, Environment, Quality of Life.* (Springer, 2019). <https://doi.org/10.1007/978-3-319-96815-5>.
42. Wang, W. M., Ding, J. L., Shu, J. W. & Chen, W. Exploration of early rice farming in China. *Quat. Int.* **227**, 22–28 (2010).
43. Kuijt, I., Finlayson, B. & MacKay, J. Pottery Neolithic landscape modification at Dhra'. *Antiquity* **81**, 106–118 (2007).
44. Gibson, S. The archaeology of agricultural terraces in the Mediterranean zone of the southern Levant and the use of the optical-stimulated luminescence (OSL) dating method. in *Soils and Sediments as Archives of Landscape Change. Geoarchaeology and Landscape Change in the Subtropics and Tropics* (eds Lucke, B., Bäuml, R. & Schmidt, M.) 295–314 (Vorstand der Fränkischen Geographischen Gesellschaft, 2015).
45. Moore, A., Mendišić, M., Smith, J., Zaninović, J. & Podrug, E. Project »Early Farming in Dalmatia«: Pokrovnik 2006 (Preliminary results). *Vjesn. Arheol. muzeja u Zagreb.* **40**, 25–34 (2007).
46. Ažman Momirski, L. Adapted slopes. *Proy. Progreso, Arqut.* **21**, 20–33 (2019).
47. Countryman, J. R. Agricultural terracing and landscape history at Monte Pallano, Abruzzo, Italy. *Honor. Pap.* **349**, (2012).
48. Denevan, W. M. *Cultivated Landscapes of Native Amazonia and the Andes* (Oxford University Press, 2001).
49. Wei, W. *et al.* Global synthesis of the classifications, distributions, benefits and issues of terracing. *Earth-Sci. Rev.* **159**, 388–403 (2016).
50. Ackermann, O., Zhevelev, H. M. & Svoray, T. Agricultural systems and terrace pattern distribution and preservation along climatic gradient: From sub-humid mediterranean to arid conditions. *Quat. Int.* **502**, 319–326 (2019).
51. Iiyama, N., Kamada, M. & Nakagoshi, N. Ecological and social evaluation of landscape in a rural area with terraced paddies in southwestern Japan. *Landsc. Urban Plan.* **73**, 60–71 (2005).
52. Goodman-Elgar, M. Evaluating soil resilience in long-term cultivation: A study of pre-Columbian terraces from the Paca Valley, Peru. *J. Archaeol. Sci.* **35**, 3072–3086 (2008).
53. *Terraced landscapes. Commemorating seventy years of the Anton Melik Geographical Institute ZRC SAZU.* (Založba ZRC, 2017).
54. Tarolli, P., Preti, F. & Romano, N. Terraced landscapes: From an old best practice to a potential hazard for soil degradation due to land abandonment. *Anthropocene* **6**, 10–25 (2014).
55. Turner, S., Bolòs, J. & Kinnaird, T. Changes and continuities in a Mediterranean landscape: A new interdisciplinary approach to understanding historic character in western Catalonia. *Landsc. Res.* **43**, 922–938 (2018).
56. Stanchi, S., Freppaz, M., Agnelli, A., Reinsch, T. & Zanini, E. Properties, best management practices and conservation of terraced soils in Southern Europe (from Mediterranean areas to the Alps): A review. *Quat. Int.* **265**, 90–100 (2012).
57. Ažman Momirski, L. Slovenian terraced landscapes. In *World Terraced Landscapes: History, Environment, Quality of Life* (eds Varotto, M. *et al.*) 45–62 (Springer, 2019). https://doi.org/10.1007/978-3-319-96815-5_4.
58. Kizos, T., Dalaka, A. & Petanidou, T. Farmers' attitudes and landscape change: Evidence from the abandonment of terraced cultivations on Lesvos, Greece. *Agric. Human Values* **27**, 199–212 (2010).
59. Ažman Momirski, L. & Kladnik, D. Terraced landscapes in Slovenia. *Acta Geogr. Slov.* **49**, 7–37 (2009).
60. Zupanc, V., Pintar, M. & Podgornik, M. Olive production on cultivated terraces in northern Istria. *Ann. Anal. za istrske mediteranske študije. Ann. di Stud. istriani e Mediterr. Ann. Istriana Mediterr. Stud.* **28**, 795–810 (2018).
61. Blanc, J.-F. Landscape typology of french agrarian terraces. In *World Terraced Landscapes: History, Environment, Quality of Life* (eds Varotto, M. *et al.*) 63–77 (Springer, 2019). https://doi.org/10.1007/978-3-319-96815-5_4.
62. Nunes, J. P. *et al.* Hydrological and erosion processes in terraced fields: Observations from a humid Mediterranean region in Northern Portugal. *L. Degrad. Dev.* **29**, 596–606 (2018).
63. Asins-Velis, S. Terraced fields in Spain: Landscapes of work and beauty. In *World Terraced Landscapes: History, Environment, Quality of Life* (eds Varotto, M. *et al.*) 79–96 (Springer, 2019).
64. Krahtopoulou, A. & Frederick, C. The stratigraphic implications of long-term terrace agriculture in dynamic landscapes: Polycyclic terracing from Kythera Island, Greece. *Geoarchaeology An Int. J.* **23**, 550–585 (2008).
65. Krüger, R. *Typologie des Waldhufendorfes nach Einzelformen und deren Verbreitungsmustern.* (Göttinger Geographische Abhandlungen, 1967).
66. Born, M. Acker- und Flurformen des Mittelalters nach Untersuchungen von Flurwüstungen. in *Untersuchungen zur eisenzeitlichen und frühmittelalterlichen Flur in Mitteleuropa und ihrer Nutzung I* (eds Beck, H., Denecke, D. & Jankuhn, H.) 310–337 (Abhandlungen der Akademie der Wissenschaften in Göttingen, Philologisch-Historische Klasse, 1979).
67. Denecke, D. Zur Terminologie ur- und frühgeschichtlicher Flurparzellierungen und Flurbegrenzungen sowie im Gelände ausgeprägter Flurrelikte, Grundzüge eines terminologischen Schemas. in *Untersuchungen zur eisenzeitlichen und frühmittelalterlichen Flur in Mitteleuropa und ihrer Nutzung I* (eds Beck, H., Denecke, D. & Jankuhn, H.) 410–440 (Abhandlungen der Akademie der Wissenschaften in Göttingen, Philologisch-Historische Klasse, 1979).
68. Sperling, W. Formen, Typen und Genese des Platzdorfes in den böhmischen Ländern: Beiträge zur Siedlungsgeographie Ostmitteleuropas. *Geogr. Zeitschrift Beih.* **61**, (1982).
69. Schmucki, R., De Blois, S., Bouchard, A. & Domon, G. Spatial and temporal dynamics of hedgerows in three agricultural landscapes of Southern Quebec, Canada. *Environ. Manage.* **30**, 0651–0664 (2002).
70. Baran-Zgłobicka, B. & Zgłobicki, W. Mosaic landscapes of SE Poland: Should we preserve them?. *Agroforest Syst* **85**, 351–365 (2012).
71. Špulerová, J., Drábová, M. & Lieskovský, J. Traditional agricultural landscape and Their Management in less Favoured areas in Slovakia. *Ekológia (Bratislava)* **35**, 1–12 (2016).
72. Dyer, C. *An Age of Transition? Economy and Society in England in the Later Middle Ages* (Clarendon Press, 2005).
73. Renes, H. Landscape history and archaeology of open fields in Europe. In *Ruralia X: Agrarian Technology in the Medieval Landscape* (ed. Klápště, J.) (Brepols Publishers n.v., 2016).

74. Dreslerová, D. Fields in Prehistoric Bohemia: Fact and Fiction. In *Agricultural and Pastoral Landscapes in Pre-industrial Society* (eds Retamero, F. et al.) 109–124 (Oxbow Books, 2016).
75. Brown, A., Walsh, K., Fallu, D., Cucchiari, S. & Tarolli, P. European agricultural terraces and lynchets: From archaeological theory to heritage management. *World Archaeol.* **52**, 566–588 (2021).
76. Andlar, G., Šrajer, F. & Trojanović, A. Classifying the mediterranean terraced landscape: The case of Adriatic Croatia. *Acta Geogr. Slov.* **57**, 111–129 (2017).
77. Lasanta, T., Errea, M. P. & Nadal-Romero, E. Traditional Agrarian landscape in the Mediterranean mountains: A regional and local factor analysis in the central Spanish Pyrenees. *L. Degrad. Dev.* **28**, 1626–1640 (2017).
78. Chen, D., Wei, W., Daryanto, S. & Tarolli, P. Does terracing enhance soil organic carbon sequestration? A national-scale data analysis in China. *Sci. Total Environ.* **721**, 137751 (2020).
79. Löw, J. & Michal, I. *Krajinný ráz [Landscape Character]*. (Lesnická práce, s.r.o., 2003).
80. Láznicka, Z. *Typy venkovského osídlení na Moravě*. (Odbor čes. společnosti zeměpisné, 1946).
81. Černý, E. *Zaniklé středověké osady a jejich pluziny. Metodika historickogeografického výzkumu v oblasti Dražanské vrchoviny*. (Academia, 1979).
82. Land Survey Office. Archive [web application]. *Central Archive of Surveying and Cadastre* <https://ags.cuzk.cz/archiv/#en> (2020).
83. Žemlička, J. *Království v pohybu. Kolonizace, města a stříbro v závěru přemyslovské epochy [Kingdom in motion. Colonization, towns and silver at the end of the Přemyslid epoch]*. (Nakladatelství Lidové noviny, 2014).
84. Demján, P. & Dreslerová, D. Modelling distribution of archaeological settlement evidence based on heterogeneous spatial and temporal data. *J. Archaeol. Sci.* **69**, 100–109 (2016).
85. Vermouzek, R. Pluzina jako datovací pomůcka [Die Flurtrassen als Datationshilfsmittel]. *Archaeol. Hist.* **7**, 265–276 (1982).
86. Klápště, J. *The Czech Lands in Medieval Transformation*. (Brill, 2012).
87. Čulíková, L. *Nedestruktivní výzkum polních systémů [Non-destructive research of field systems]*. (University of West Bohemia in Pilsen, 2013).
88. Sadravetová, M. Nástin středověké a novověké kolonizace Vimperska [The Outline of Medieval and Postmedieval Colonization of the Vimperk Region]. *Hist. geogr.* **41**, 107–130 (2015).
89. Fanta, V., Zouhar, J., Beneš, J., Bumerl, J. & Sklenicka, P. How old are the towns and villages in Central Europe? Archaeological data reveal the size of bias in dating obtained from traditional historical sources. *J. Archaeol. Sci.* **113**, (2020).
90. Fanta, V., Zouhar, J., Beneš, J., Bumerl, J. & Sklenicka, P. On the time-lag in the dating of historical settlements: Reply to Kolář and Szabó (2021). *J. Archaeol. Sci.* **132**, 105391 (2021).
91. Dohnal, M. Regulace vesnického osídlení a krajiny ve středověké a novověké severní Evropě. *Český lid. Etnol. časopis = Ethnol. J.* **96**, 279–308 (2009).
92. Sádlo, J., Pokorný, P., Hájek, P., Dreslerová, D. & Čilek, V. *Krajina a revoluce: Významné přelomy ve vývoji kulturní krajiny Českých zemí*. (Malá Skála, 2008).
93. Thomas, G., McDonnell, G., Merkel, J. & Marshall, P. Technology, ritual and Anglo-Saxon agriculture: The biography of a plough coulter from Lyminge, Kent. *Antiquity* **90**, 742–758 (2016).
94. Banham, D. & Faith, R. *Anglo-Saxon Farms and Farming. Anglo-Saxon Farms and Farming* (Oxford University Press, 2014). <https://doi.org/10.1093/ACPROF/OSO/9780199207947.001.0001>.
95. Klír, T. *Osídlení zemědělsky marginálních půd v mladším středověku a raném novověku* (Univerzita Karlova, 2008).
96. Němčenko, N. *Dějiny pozemkových úprav II*. (České vysoké učení technické v Praze, 1970).
97. Žemlička, J. Osídlení Zbraslavka od 10. do počátku 15. století. *Památky archeol.* **65**, 419–465 (1974).
98. Boháč, Z. Geneze sídla a pluziny jako pramen k dějinám osídlení. *Hist. geogr.* **25**, 7–52 (1986).
99. Fanta, V. The use of long-term historical data in exploring landscape-ecological issues. (Czech University of Life Sciences Prague [PhD thesis], 2019).
100. Lucke, B. et al. Bodenentwicklung im Kontext der historischen Landnutzung und Geologie der Gemarkung Fatschenbrunn im Steigerwald. *Mitteilungen der Fränkischen Geogr. Gesellschaft* **61**(62), 71–82 (2016).
101. Kladnik, D. & Perko, D. Terraced landscapes in Slovenia. in *Terraced Landscapes. Commemorating seventy years of the Anthon Melik Geographical Institute ZRC SAZU* (eds Perko, D., Ciglič, R., Geršič, M. & Kladnik, D.) (Založba ZRC, 2017).
102. Ažman Momirski, L. & Berčić, T. Ignored regions: Slovenian terraced landscapes. *Ann. Ann. Istrian Mediterr. Stud. Ser. Hist. Sociol.* **26**, 399–418 (2016).
103. Žemlička, J. *Vývoj osídlení dolního Poohří a Českého středohoří do 14. století*. (Academia, 1980).
104. *A Companion to Social Archaeology*. (Blackwell Publishing Ltd, 2007). <https://doi.org/10.1002/9780470693605>.
105. Oosthuizen, S. M. *Tradition and Transformation in Anglo-Saxon England: Archaeology, Common Rights and Landscape* (Bloomsbury Academic, 2013). <https://doi.org/10.5040/9781472555861>.
106. Born, M. *Geographie der ländlichen Siedlungen 1. Die Genese der Siedlungsformen in Mitteleuropa*. (B.G. Teubner, 1977).
107. Hopcroft, R. L. *Regions, Institutions, and Agrarian Change in European History* (University of Michigan Press, 1999). <https://doi.org/10.3998/MPUB.16130>.
108. Fanta, V., Janečková, K. & Černý Pixová, K. Identifikace a ochrana dochovaných pozůstatků historických pluzin: zpráva o projektu NAKI [Identification and preservation of historic field patterns: National Cultural Identity project report]. *Pozem. úpravy* **29**, 3–6 (2021).
109. Bonardi, L. & Mastrovito, D. Italian terraced vineyards: A geographical overview. *Pirineos* **175**, (2020).
110. Špulerová, J. et al. Inventory and classification of historical structures of the agricultural landscape in Slovakia. *Ekológia (Bratislava)* **30**, 157–170 (2011).
111. Kladnik, D., Kruse, A. & Komac, B. Terraced landscapes: An increasingly prominent cultural landscape type. *Acta Geogr. Slov.* **57**, 73–81 (2017).
112. Semotanová, E. *Historická geografie Českých zemí [Historical Geography of the Czech Lands]*. (Historický ústav AV ČR, 2006).
113. Hauserová, M. & Poláková, J. *Pomůcka pro používání základních historických map*. (Ústav památkové péče FA ČVUT, 2015).
114. Klápště, J. Středověké osídlení Černokostelecka. *Památky Archeol.* **69**, 423–475 (1978).
115. Nekuda, V. *Pfaffenschlag: zaniklá středověká ves u Slavonic: příspěvek k dějinám středověké vesnice*. (Blok a Moravské Muzeum, 1975).
116. Smetánka, Z. *Život středověké vesnice: Zaniklá Svídna*. (Academia, 1988).
117. Belcredi, L. *Bystřec: o založení, životě a zániku středověké vsi: archeologický výzkum zaniklé středověké vsi na Dražanské vrchovině 1975–2005*. (Muzejní a vlastivědná společnost v Brně: Moravské zemské muzeum, 2006).
118. Smetánka, Z. & Klápště, J. Geodeticko-topografický průzkum zaniklých středověkých vsí na Černokostelecku [Geodetical-topographical survey of deserted medieval villages in the Kostelec-nad-Černými-lesy region]. *Památky Archeol.* **72**, 416–458 (1981).
119. *Archeologie a letecké laserové skenování krajiny [Archaeology and airborne laser scanning of the landscape]*. (Katedra archeologie, Západočeská univerzita v Plzni, 2013).
120. Houfková, P. et al. The dynamics of a non-forested stand in the Krušné Mts.: The effect of a short-lived medieval village on the local environment. *Veg. Hist. Archaeobot.* **28**, 607–621 (2019).
121. Beneš, J., Hrubý, P., Michálek, J. & Parkman, M. Kamenná hrazení na Hořejším vrchu a vrchu Kokovci u Vlachova Březí. *Zlatá stezka Sborník Pr. Muz.* **6**, 271–296 (1999).

122. Zimová, K. *et al.* Analýza vývoje plůžiny zaniklé obce Malonín na Prachaticku. *Acta Pruhoniciana* **104**, 27–37 (2013).
123. Horák, J. & Klír, T. Pedogenesis, Pedochemistry and the Functional Structure of the Waldhufendorf Field System of the Deserted Medieval Village Spindelbach, the Czech Republic. *Interdiscip. Archaeol. Nat. Sci. Archaeol.* **8**, (2017).
124. Horák, J., Janovský, M., Hejzman, M., Šmejda, L. & Klír, T. Soil geochemistry of medieval arable fields in Lovětín near Třešň, Czech Republic. *CATENA* **162**, 14–22 (2018).
125. Černý, E. *Výsledky výzkumu zaniklých středověkých osad a jejich plůžin. Historicko-geografická studie v regionu Dražanské vrchoviny [The results of research on deserted medieval villages and their field systems. (...)].* (Muzejní a vlastivědná společnost, 1992).
126. Molnářová, K. Hedgerow-defined medieval field patterns in the Czech Republic and their conservation: A literature review. *J. Landsc. Stud.* **1**, 27–47 (2008).
127. Čílek, V. *et al.* *Vstoupit do krajiny. O přírodě a paměti středních Čech [Enter the Landscape. The Nature and Memory of Central Bohemia]*. (Dokořán, 2004).
128. Chuman, T. & Romportl, D. Multivariate classification analysis of cultural landscapes: An example from the Czech Republic. *Landsc. Urban Plan.* **98**, 200–209 (2010).
129. Löw, J. & Novák, J. Typologické členění krajiny České republiky: Typological Subdivision of Landscapes in the Czech Republic. *Urban. a územní Rozv.* **11**, 19–23 (2008).
130. Pešta, J. *Encyklopedie českých vesnic. Vesnické památkové rezervace, zóny a ostatní památkově hodnotná vesnická sídla v Čechách. Díl I. Střední Čechy a Praha.* (Libri, 2003).
131. Bartlett, R. *The Making of Europe. Conquest, Colonization and Cultural Change 950–1350.* (Penguin Books, 1994).
132. Kuča, K. *Města a městečka v Čechách, na Moravě a ve Slezsku. III. díl: Kolín–Mí [Towns and townships in Bohemia, Moravia and Silesia. Part III.: Kolín–Mí]*. (Libri, 1998).
133. Vařeka, P., Holata, L., Rožmberský, P. & Schejbalová, Z. Středověké osídlení Rokycanska a problematika zaniklých vsí: The Mediaeval Settlement of the Rokycansko Region and the Issue of Deserted Villages. *Archaeol. Hist.* **36**, 319–342 (2011).
134. Odložilík, O., Zeman, Z. A. B., Bradley, J. F. N., Hauner, M. & Wiskemann, E. Czechoslovak history. *Encyclopædia Britannica* <https://www.britannica.com/topic/Czechoslovak-history> (2016).
135. Kuna, M. *et al.* *Nedestruktivní archeologie. Teorie, metody a cíle: Non-destructive archaeology. Theory, methods and goals.* (Academia, 2004).
136. Munck, T. *Seventeenth-Century Europe 1598–1700: State, Conflict and the Social Order in Europe* (Palgrave Macmillan, 2005).
137. Fanta, V., Šálek, M., Zouhar, J., Sklenicka, P. & Storch, D. Equilibrium dynamics of European pre-industrial populations: The evidence of carrying capacity in human agricultural societies. *Proc. R. Soc. B Biol. Sci.* **285**, 20172500 (2018).
138. Kirsten, E., Buchholz, E. W. & Köllmann, W. *Raum und Bevölkerung in der Weltgeschichte [Space and Population in the World's History]*. (A. G. Ploetz – Verlag, 1965).
139. Lederer, D. The Myth of the All-Destructive War: Afterthoughts on German Suffering, 1618–1648. *Ger. Hist.* **29**, 380–403 (2011).
140. Fialová, L. *et al.* *Dějiny obyvatelstva českých zemí [Population History of the Czech Lands]*. (Mladá fronta, 1998).
141. Maršíková, M. & Maršík, Z. *Dějiny zeměměřičství a pozemkových úprav v Čechách a na Moravě v kontextu světového vývoje.* (Libri, 2007).
142. Matoušek, V. *Čechy krásné, Čechy mé. Proměny krajiny Čech v době industriální [My beautiful Bohemia. Changes of the Czech landscape in industrial era]*. (Krigl, 2010).
143. Hájek, P. *Jde pevně kupředu naše zem. Krajina českých zemí v období socialismu 1948–1989.* (Malá Skála, 2008).
144. Bičík, I., Jeleček, L. & Štěpánek, V. Land-use changes and their social driving forces in Czechia in the 19th and 20th centuries. *Land Use Policy* **18**, 65–73 (2001).
145. Bičík, I. *et al.* Land Use/Cover Changes in Czechia over the Past 150 Years - An Overview. in *Land Use/Cover Changes in Selected Regions in the World* (eds. Himiyama, Y., Mather, A., Bičík, I. & Milanova, E. V.) vol. 1 29–39 (International Geographical Union Commission on Land Use and Land Cover Change (IGU-LUCC), 2001).
146. Skaloš, J. *et al.* Using old military survey maps and orthophotograph maps to analyse long-term land cover changes: Case study (Czech Republic). *Appl. Geogr.* **31**, 426–438 (2011).
147. Vaishar, A., Vavrouchová, H., Lešková, A. & Peřínková, V. Depopulation and Extinction of Villages in Moravia and the Czech Part of Silesia since World War II. *Land* **10**, 333 (2021).
148. Zelinka, V., Zacharová, J. & Skaloš, J. Analysis of spatiotemporal changes of agricultural land after the Second World War in Czechia. *Sci. Rep.* **11**, 12655 (2021).
149. Frielinghaus, M. & Schmidt, R. Historische landschaftsanalyse und abgeleitete aktuelle entwicklungsziele. *Arch. Agron. Soil Sci.* **47**, 73–91 (2001).
150. Lipský, Z. *Sledování změn v kulturní krajině [Monitoring of changes in cultural landscapes]*. (Česká zemědělská univerzita, 2000).
151. Institute of Archaeology of the Czech Academy of Sciences Prague. Archeologická databáze Čech [Archaeological Database of Bohemia]. *Archeologická mapa ČR [Archaeological Map of the Czech Republic]* <http://archeologickamapa.cz/downloads/ADC2014.mdb> (2013).
152. Wikipedia. Franziszeischer Kataster. *Wikipedia, Die freie Enzyklopädie.* https://de.wikipedia.org/w/index.php?title=Franziszeischer_Kataster&oldid=208057787 (2021).
153. Fanta, V. & Rákosníková, A. Lidová architektura českých vesnic v Banátu. in *Přenesená krajina. Český venkov v rumunském Banátu* (ed. Kovář, P.) 67–113 (Academia, 2019).
154. Hendrychová, M. & Kabrna, M. An analysis of 200-year-long changes in a landscape affected by large-scale surface coal mining: History, present and future. *Appl. Geogr.* **74**, 151–159 (2016).
155. Pavelková, R. *et al.* Historical ponds of the Czech Republic: An example of the interpretation of historic maps. *J. Maps* **12**, 551–559 (2016).
156. Pešta, J. *Plošný průzkum lidové architektury a venkovských sídel.* (Národní památkový ústav, územní odborné pracoviště středních Čech v Praze, 2014).
157. Máčel, O. *Základní problematika urbanistické struktury vesnice v Čechách a na Moravě I a II.* (1955).
158. GISAT. SRTM DEM [data]. "Data provided and processed by © GISAT (2007)". http://www.gisat.cz/data/dem/SRTM_DEM_CZ_KR_100.zip (2007).
159. Czech Geological Survey. Geological map of the Czech Republic 1 : 500 000. *Map services based on Esri technology* <https://mapy.geology.cz/geocr500/> (2013).
160. State Land Office. BPEJ [Soil units]. *Celostátní databáze BPEJ [National database of soil units]* https://www.spucr.cz/frontend/webroot/uploads/files/2019/04/bpej_201904019035.zip (2021).
161. Bečvářová, V., Vašek, P. & Vaníček, F. *Bonitace čs. zemědělských půd a směry jejich využití. 4. díl. Zařazení zemědělských podniků a katastrů do produkčně-ekonomických skupin (PES) [Rating of Czechoslovak Agricultural Soils and Trends in their Use. 4th part]*. (Federální ministerstvo zemědělství a výživy: Ministerstvo zemědělství ČSR: Ministerstvo poľnohospodárstva a výživy SSR, 1988).
162. *Atlas československých dějin [Atlas of Czechoslovak history]*. (Ústřední správa geodézie a kartografie: Historický ústav Československé akademie věd: Kartografický a reprodukční ústav v Praze, 1965).
163. *Atlas krajiny české republiky: Landscape Atlas of the Czech Republic.* (Ministerstvo životního prostředí ČR: Výzkumný ústav Silva Taroucy pro krajinu a okrasné zahradnictví, v. v. i., 2009).

164. Arcdata Praha. ArcCR® 500, version 3.3 [ESRI database]. <http://download.arcdata.cz/data/ArcCR500-3.3-windows-installer.exe> (2016).
165. Kuča, K. Půdorysné struktury [Layout compositions]. *CZ_Retro* <https://geoportal.npu.cz/webappbuilder/apps/38/> (2017).
166. *Akademický atlas českých dějin [Academic atlas of Czech history]*. (Academia, 2014).
167. Klír, T. Osídlení horských oblastí Čech ve středověku a raném novověku: východiska interdisciplinárního výzkumu: Settlement in Mountainous Areas of Bohemia in the Middle Ages and the Early Modern Age: Points of Departure for Interdisciplinary Research. *Archaeol. Hist.* **35**, 373–391 (2009).
168. QGIS Development Team. QGIS Geographic Information System [software]. (2017).
169. Papke, L. E. & Wooldridge, J. M. Econometric methods for fractional response variables with an application to 401(k) plan participation rates. *J. Appl. Econom.* **11**, 619–632 (1996).
170. Buis, M. L. FMLOGIT: Stata module fitting a fractional multinomial logit model by quasi maximum likelihood. *Statistical Software Components* S456976 (2008).
171. Burnham, K. P. *Multimodel inference: Understanding AIC relative variable importance values*. <https://sites.warnercnr.colostate.edu/kenburnham/wp-content/uploads/sites/25/2016/08/VARIMP.pdf> (2015).
172. Symonds, M. R. E. & Moussalli, A. A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's information criterion. *Behav. Ecol. Sociobiol.* **65**, 13–21 (2011).
173. Luchman, J. N. MIINC: Stata module to conduct multi-model inference using information criteria. *Statistical Software Components* S457828 (2014).
174. Barbieri, M. M. & Berger, J. O. Optimal predictive model selection. *Ann. Stat.* **32**, 870–897 (2004).
175. Jann, B. Making regression tables from stored estimates. *Stata J.* **5**, 288–308 (2005).
176. Conrad, O. *et al.* System for Automated Geoscientific Analyses (SAGA) v. 2.1.4 [software]. *Geosci. Model Dev.* 1991–2007 (2015). 10.5194/gmd-8-1991-2015.
177. *Encyclopedia of Biodiversity*. (Academic Press, 2013).
178. Domagal-Goldman, S. D. *et al.* The Astrobiology Primer v2.0. *Astrobiology* **16**, 561–653 (2016).
179. Merino, N. *et al.* Living at the extremes: Extremophiles and the limits of life in a planetary context. *Front. Microbiol.* **10**, 780 (2019).
180. Niederberger, T. Extremophile. *Encyclopædia Britannica* <https://www.britannica.com/science/extremophile> (2021).
181. Beranová, M. & Kubačák, A. *Dějiny zemědělství v Čechách a na Moravě [The history of agriculture in Bohemia and Moravia]*. (Libri, 2010).
182. Dohnal, M. *Vesnická sídla a kulturní krajina na Táborsku v 15.–19. století: Villages and cultural landscapes in the region of Tábor from the 15th to 19th century*. (Etnologický ústav Akademie věd České republiky, 2006).
183. Krenzlín, A. & Reusch, L. *Die Entstehung der Gewinnflur nach Untersuchungen im nördlichen Unterfranken*. (Kramer, 1961).
184. Dohnal, M. Late medieval and early modern modifications of regular open-field systems in Bohemia. *Bohemia A J. Hist. Civilis. East Cent. Eur.* **56**, 35–61 (2016).
185. Scharlau, K. Gewinnflurenforschung in Hessen. *Vor- und Forschungen Die Anfänge der Landgemeinde und ihr Wes.* **17**, 29–51 (1964).
186. Eyre, S. R. The Curving Plough-strip and its Historical Implications. *Agric. Hist. Rev.* **3**, 80–94 (1955).
187. Robinson, M. L. & Strauss, S. Y. Generalists are more specialized in low-resource habitats, increasing stability of ecological network structure. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 2043–2048 (2020).
188. Dohnal, M. & Škabrada, J. Utváření vesnických normových půdorysů na Olomoucku ve středověku a novověku. *Český lid. Etnol. časopis Ethnol. J.* **102**, 199–232 (2015).
189. Göransson, S. Field and village on the Island of Öland: A study of the genetic compound of an east Swedish rural landscape. *Geogr. Ann.* **40**, 101–158 (1958).
190. Göransson, S. Solskifte: The Definition of a Confused Concept. in *Fields, Farms and Settlement in Europe* (eds. Buchanan, R. H., Butlin, R. A. & McCourt, D.) 22–37 (Cultra Manor, 1976).
191. Helmfrid, S. Östergötland 'vastianstang': Studien über die altere Agrarlandschaft und ihre Genese. *Geogr. Ann.* **44**, 1–277 (1962).
192. Semotanová, E. Vogtova veduta Lysé nad Labem: 'naučná stezka' Šporkovou barokní krajinou. in *Od knížat ke králům: Sborník u příležitosti 60. narozenin Josefa Žemličky* (eds. Doležalová, E., Šimůnek, R., Dvořáčková-Malá, D. & Pořízka, A.) 494–514 (Nakladatelství Lidové noviny, 2007).
193. Kaše, J. & Kotlík, P. *Braunův Betlém. Drama krajiny a umění v proměnách času*. (Paseka, 1999).
194. Charvátová, K. Pozemková držba českých cisterciáckých klášterů 1142–1420: Landed property of the Cistercian houses of Bohemia 1142–1420. *Architectus* **3**, 3–16 (2013).
195. Bahenský, F., Dohnal, M., Vařeka, J. & Woitsch, J. *Etnografický atlas Čech, Moravy a Slezska VII. Lidové stavitelství*. (Etnologický ústav Akademie věd České republiky, v. v. i., 2015).
196. Zhang, Y. *et al.* Patterns in pottery use reveal different adaptive strategies between lower and higher altitude regions on the Tibetan Plateau: Chemical evidence from pottery residues. *J. Archaeol. Sci.* **138**, 105544 (2022).

Acknowledgements

This research was supported by IGA projects No. 2020B0042 “Rychlost zániku krajinných prvků coby nositelů paměti kulturní krajiny” and No. 2021B0046 “Proměna krajinné mikrostruktury v Evropě” of the Faculty of Environmental Sciences, Czech University of Life Sciences Prague. Further support was received from NAKI II project No. DG18P02OVV060 “Identifikace a ochrana dochovaných pozůstatků historických plůžin” of the Ministry of Culture of the Czech Republic, and from Prostředí pro život / Environment for Life project No. SS02030018 “Centrum pro krajinu a biodiverzitu” of the Technology Agency of the Czech Republic. We thank David Kalhous and Dagmar Dreslerová for their fruitful comments. We also thank Markéta Šantrůčková for her assistance with translations of German terms, Radana “Radouch” Kavková for her explanation of the geological context and Jiří “Bumča” Bumerl for his help with the archaeological data. We thank Kateřina Fantová and Robin Healey for language assistance.

Author contributions

V.F. and V.R. collected the data, V.F. and J.Z. analysed the data, V.F. prepared the figures, and V.F., J.B., J.Z., L.Š., K.J.M. and P.S. wrote the paper.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-12612-8>.

Correspondence and requests for materials should be addressed to V.F.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022